



ST2008022A

2008 Aerial Imagery With Soil Survey



ST2008022A

Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics





Trial Type	Headline Fungicide
Application Method	In Furrow
Crop Rotation	Corn Following Corn (4 Years)
Hybrid/Variety	NK Brand N60-B4
Application Date/Crop Stage	4/30/2008 - Planting
County	Black Hawk

FUNGICIDE TREATMENT
Yield Data Points
● Headline
● Untreated
● Non-Trial



ST2008022A

Trial Grain Yield Map With Soil Survey



ST2008022A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Dickinson Fine Sandy Loam, 0 To 2 Percent Slopes	175	3.9	4.0	188.0	193.0	-5.0
Saude Loam, 0 To 2 Percent Slopes	177	5.5	6.4	183.3	178.6	4.7
Waukee Loam, 0 To 2 Percent Slopes	178	11.0	11.5	211.0	207.9	3.1
Flagler Sandy Loam, 0 To 2 Percent Slopes	284	10.1	11.0	185.2	184.2	1.0
Chelsea Loamy Fine Sand, 9 To 14 Percent Slopes	63D	3.1	4.4	198.2	195.2	3.0
Wiota Silty Clay Loam, 0 To 2 Percent Slopes	7	14.3	14.7	199.6	192.6	7.0
YIELD AVERAGE FOR TRIAL				196.3	192.7	3.6

