



ST2008022B

2008 Aerial Imagery With Soil Survey



ST2008022B

Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics



Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Corn Following Corn (4 Years)
Hybrid/Variety	NK Brand N60-B4
Application Date/Crop Stage	7/26/2008 - R1 Silking
County	Black Hawk

FUNGICIDE TREATMENT



Yield Data Points

- Headline
- Untreated
- Non-Trial



ST2008022B

Trial Grain Yield Map With Soil Survey



ST2008022B

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Dickinson Fine Sandy Loam, 0 To 2 Percent Slopes	175	3.9	2.4	192.8	197.9	-5.1
Saude Loam, 0 To 2 Percent Slopes	177	12.9	2.1	182.0	182.2	-0.2
Waukee Loam, 0 To 2 Percent Slopes	178	2.0	18.2	197.9	210.1	-12.2
Flagler Sandy Loam, 0 To 2 Percent Slopes	284	20.0	2.5	183.8	196.7	-12.9
Sparta Loamy Fine Sand, 9 To 14 Percent Slopes	41D	0.0	0.2	0.0	225.9	0.0
Chelsea Loamy Fine Sand, 9 To 14 Percent Slopes	63D	0.0	0.3	0.0	212.5	0.0
Wiota Silty Clay Loam, 0 To 2 Percent Slopes	7	10.9	24.6	185.1	197.1	-12.0
YIELD AVERAGE FOR TRIAL						
				184.9	201.4	-16.5

