



ST2008239A

2008 Aerial Imagery With Soil Survey



ST2008239A

Treatment Type and Location of Trial Within Field






Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Corn Following Corn (2 Years)
Hybrid/Variety	NK Brand N48G
Application Date/Crop Stage	7/31/2008 - R1 Silking
County	Black Hawk

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008239A

Trial Grain Yield Map With Soil Survey



ST2008239A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Marshan Clay Loam, 24 To 40 Inches To Sand And Gravel, 0 To 2 Percent Slopes	1152	10.1	11.1	189.0	181.7	7.3
Dickinson Fine Sandy Loam, 2 To 5 Percent Slopes	175B	19.9	16.5	176.9	172.2	4.7
Oran Loam, 1 To 3 Percent Slopes	471	2.8	1.8	167.5	148.2	19.3
Chelsea Loamy Fine Sand, 5 To 9 Percent Slopes	63C	3.4	2.8	133.9	124.3	9.6
Chelsea Loamy Fine Sand, 9 To 14 Percent Slopes	63D	1.8	1.6	147.0	141.5	5.5
Wiota Silty Clay Loam, 0 To 2 Percent Slopes	7	12.5	15.9	186.8	178.4	8.4
YIELD AVERAGE FOR TRIAL				177.3	171.5	5.8

