



ST2008250A

2008 Aerial Imagery With Soil Survey



ST2008250A

Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics



Trial Type	Stratego Fungicide (2 Pass)
Application Method	Ground Applied
Crop Rotation	Corn Following Soybeans
Hybrid/Variety	NK Brand N48-B1
Application Date/Crop Stage	8/2/2008 - R1 Silking
County	Black Hawk

FUNGICIDE TREATMENT	
Yield Data Points	
●	Stratego
●	Untreated
●	Non-Trial



ST2008250A

Trial Grain Yield Map With Soil Survey



ST2008250A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Stratego	Untreated	Stratego	Untreated	
Dickinson Fine Sandy Loam, 2 To 5 Percent Slopes	175B	27.4	30.0	158.4	161.1	-2.7
Sparta Loamy Fine Sand, 2 To 5 Percent Slopes	41B	13.3	12.1	113.3	119.7	-6.4
Sparta Loamy Fine Sand, 5 To 9 Percent Slopes	41C	2.9	2.3	80.7	80.8	-0.1
Oran Loam, 1 To 3 Percent Slopes	471	4.4	4.0	119.9	141.3	-21.4
Chelsea Loamy Fine Sand, 2 To 5 Percent Slopes	63B	1.8	1.9	125.1	130.0	-4.9
YIELD AVERAGE FOR TRIAL				137.2	144.8	-7.6

