




ST2008233A

# 2008 Aerial Imagery With Soil Survey



ST2008233A

## Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics

<b>Trial Type</b>	Headline Fungicide
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Corn Following Soybeans
<b>Hybrid/Variety</b>	NK Brand NK 67-W2
<b>Application Date/Crop Stage</b>	7/28/2008 - R1 Silking
<b>County</b>	Bremer

### FUNGICIDE TREATMENT



Yield Data Points

- Headline
- Untreated
- Non-Trial



ST2008233A

# Trial Grain Yield Map With Soil Survey



ST2008233A

## Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Atkinson	AtC	16.0	13.5	199.0	196.5	2.5
Cresco	CrC	1.6	0.7	196.5	198.8	-2.3
Dinsdale	DsB	26.6	19.7	214.1	206.7	7.4
Rockton	RkD	12.9	8.0	173.5	186.2	-12.7
Sogn Soils	SoD	0.2	0.0	202.9	0.0	0.0
Terril	TxB	0.7	0.2	150.8	155.0	-4.2
<b>YIELD AVERAGE FOR TRIAL</b>				<b>199.6</b>	<b>199.2</b>	<b>0.4</b>

