



ST2008258A

# 2008 Aerial Imagery With Soil Survey



ST2008258A

# Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics






<b>Trial Type</b>	Headline Fungicide
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Soybeans Following Corn
<b>Hybrid/Variety</b>	FS/Growmark HS2555
<b>Application Date/Crop Stage</b>	8/8/2008 - R3 Beg. Pod
<b>County</b>	Bremer

<b>FUNGICIDE TREATMENT</b>	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008258A

# Trial Grain Yield Map With Soil Survey



ST2008258A

## Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Atkinson	AtB	3.9	3.7	49.0	47.1	1.9
Bassett	BeB	2.0	1.0	38.6	37.0	1.6
Dinsdale	DsB	10.6	11.3	46.3	44.0	2.3
Kenyon	KeB	4.9	2.5	42.7	38.6	4.1
Kenyon	KeC	11.7	11.2	41.7	37.8	3.9
Klinger	Kg	11.3	10.9	42.7	40.0	2.7
Terril	TxA	1.5	1.5	43.0	43.5	-0.5
Terril	TxB	3.1	2.1	42.1	40.2	1.9
Waukegan Deep	WkA	4.5	2.3	47.2	46.5	0.7
<b>YIELD AVERAGE FOR TRIAL</b>				<b>43.9</b>	<b>41.3</b>	<b>2.6</b>

