


ST2008400A


Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics



Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	Asgrow AG2606
Application Date/Crop Stage	8/10/2008 - R5 Beg. Seed
County	Buchanan

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008400A

Trial Grain Yield Map With Soil Survey



ST2008400A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Rockton Loam, 30 To 40 Inches To Limestone, 2 To 6 Percent Slopes	213B	0.5	0.5	50.1	48.5	1.6
Clyde-Floyd Complex, 1 To 4 Percent Slopes	391B	0.9	1.1	51.0	51.3	-0.3
Olin Fine Sandy Loam, 2 To 5 Percent Slopes	408B	0.8	0.0	36.7	0.0	0.0
Sparta Loamy Fine Sand, 2 To 5 Percent Slopes	41B	1.9	2.0	42.2	42.4	-0.2
Spillville Loam, 0 To 2 Percent Slopes	485	1.1	1.0	49.0	44.8	4.2
Bertram Fine Sandy Loam, 2 To 6 Percent Slopes	809B	4.1	4.5	42.8	41.1	1.7
Kenyon Loam, 2 To 5 Percent Slopes	83B	43.2	38.4	50.5	52.4	-1.9
YIELD AVERAGE FOR TRIAL				49.4	50.7	-1.3

