



ST2008288A

2008 Aerial Imagery With Soil Survey




ST2008288A

Treatment Type and Location of Trial Within Field





Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	NK Brand S17-Z7
Application Date/Crop Stage	? - ?
County	Chickasaw

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008288A

Trial Grain Yield Map With Soil Survey



ST2008288A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Bassett Loam, 2 To 5 Percent Slopes	171B	1.3	3.1	47.3	46.4	0.9
Floyd Loam, 1 To 4 Percent Slopes	198B	12.0	11.6	43.6	41.1	2.5
Readlyn Loam, 2 To 5 Percent Slopes	399B	9.0	3.4	49.4	45.8	3.6
Protivin Loam, 1 To 4 Percent Slopes	798B	7.1	6.8	46.1	38.9	7.2
Kenyon Loam, 2 To 5 Percent Slopes	83B	6.7	9.2	45.9	45.9	0.0
Kenyon Loam, 5 To 9 Percent Slopes, Moderately Eroded	83C2	5.2	6.8	49.4	48.6	0.8
Clyde Clay Loam, 0 To 3 Percent Slopes	84	8.6	9.2	48.5	45.2	3.3
YIELD AVERAGE FOR TRIAL				46.9	44.1	2.8

