



ST2008124A

2008 Aerial Imagery With Soil Survey



ST2008124A

Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics




Trial Type	Domark Fungicide
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	Pioneer Experimental
Application Date/Crop Stage	8/7/2008 - Not Provided
County	Greene

FUNGICIDE TREATMENT	
Yield Data Points	
●	Domark
●	Untreated
●	Non-Trial



ST2008124A

Trial Grain Yield Map With Soil Survey



ST2008124A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Domark	Untreated	Domark	Untreated	
Coland Silty Clay Loam, 0 To 2 Percent Slopes	135	0.5	1.1	60.9	57.9	3.0
Clarion Loam, 2 To 5 Percent Slopes	138B	6.9	1.9	62.1	59.7	2.4
Clarion Loam, 2 To 5 Percent Slopes, Moderately Eroded	138B2	7.7	9.6	59.1	59.7	-0.6
Marna Silty Clay Loam, 0 To 2 Percent Slopes	383	10.8	18.5	58.4	59.9	-1.5
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	5.5	9.2	60.0	59.1	0.9
Nicollet Loam, 1 To 3 Percent Slopes	55	0.0	0.7	0.0	54.1	0.0
Clarion-Storden Loams, 5 To 9 Percent Slopes, Moderately Eroded	638C2	9.9	10.8	56.7	57.1	-0.4
Clarion-Storden Loams, 9 To 14 Percent Slopes, Moderately Eroded	638D2	3.6	3.4	53.5	54.2	-0.7
YIELD AVERAGE FOR TRIAL				58.6	58.7	-0.1

