




ST2008130A

# 2008 Aerial Imagery With Soil Survey



ST2008130A


# Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics




<b>Trial Type</b>	Headline Fungicide
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Soybeans Following Corn
<b>Hybrid/Variety</b>	Stine S2032-4
<b>Application Date/Crop Stage</b>	7/31/2008 - Not Provided
<b>County</b>	Greene

<b>FUNGICIDE TREATMENT</b>	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008130A

# Trial Grain Yield Map With Soil Survey



ST2008130A

# Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference		
		Headline	Untreated	Headline	Untreated			
Webster Clay Loam, 0 To 2 Percent Slopes	107	14.5	13.8	44.4	39.5	4.9		
Clarion Loam, 2 To 5 Percent Slopes, Moderately Eroded	138B2	17.8	19.7	46.0	40.6	5.4		
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	4.1	3.2	43.7	40.6	3.1		
Nicollet Loam, 1 To 3 Percent Slopes	55	8.3	8.4	47.1	41.2	5.9		
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	0.1	0.2	40.8	41.4	-0.6		
Calco Silty Clay Loam, 0 To 2 Percent Slopes	733	4.5	5.2	41.1	36.9	4.2		
<b>YIELD AVERAGE FOR TRIAL</b>						<b>45.1</b>	<b>40.0</b>	<b>5.1</b>

