



ST2008160A

2008 Aerial Imagery With Soil Survey



ST2008160A

Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics

Trial Type	Domark Fungicide
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	NuTech Seed 2660
Application Date/Crop Stage	8/1/2008 - Not Provided
County	Greene

FUNGICIDE TREATMENT




Yield Data Points

- Domark
- Untreated
- Non-Trial



ST2008160A

Trial Grain Yield Map With Soil Survey



ST2008160A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Domark	Untreated	Domark	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	18.7	19.7	45.2	48.2	-3.0
Clarion Loam, 2 To 5 Percent Slopes	138B	3.6	4.4	47.7	48.3	-0.6
Clarion Loam, 2 To 5 Percent Slopes, Moderately Eroded	138B2	13.2	15.1	46.9	48.7	-1.8
Knoke Silty Clay Loam, 0 To 1 Percent Slopes	4	0.4	0.0	43.9	0.0	0.0
Nicollet Loam, 1 To 3 Percent Slopes	55	8.3	9.2	47.6	48.7	-1.1
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	2.9	3.4	43.6	45.6	-2.0
Clarion-Storden Loams, 5 To 9 Percent Slopes, Moderately Eroded	638C2	0.5	0.2	51.5	42.1	9.4
Harps Loam, 0 To 2 Percent Slopes	95	0.4	0.0	41.8	0.0	0.0
YIELD AVERAGE FOR TRIAL				46.2	48.2	-2.0

