




ST2008392A

2008 Aerial Imagery With Soil Survey



ST2008392A

Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics

Trial Type	Plant Nutrition
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	? ?
Application Date/Crop Stage	7/31/2008 - ?
County	Greene




PLANT TREATMENT

- Yield Data Points
- Monty's Plant Food
 - Untreated
 - Non-Trial



ST2008392A

Trial Grain Yield Map With Soil Survey



ST2008392A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Monty's Plant Food	Untreated	Monty's Plant Food	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	13.1	11.5	45.7	46.8	-1.1
Clarion Loam, 2 To 5 Percent Slopes	138B	5.3	5.2	46.0	47.2	-1.2
Clarion Loam, 5 To 9 Percent Slopes, Moderately Eroded	138C2	3.2	3.3	43.8	44.0	-0.2
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	2.1	2.9	49.1	49.9	-0.8
Nicollet Loam, 1 To 3 Percent Slopes	55	8.2	9.6	49.4	48.7	0.7
Clarion-Storden Loams, 5 To 9 Percent Slopes, Moderately Eroded	638C2	12.1	13.8	43.7	42.8	0.9
Salida-Storden Complex, 5 To 9 Percent Slopes, Moderately Erode	639C2	4.9	4.8	38.4	40.2	-1.8
YIELD AVERAGE FOR TRIAL				45.1	45.5	-0.4

