



ST2008061A

# 2008 Aerial Imagery With Soil Survey



ST2008061A

# Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics

<b>Trial Type</b>	Extreme
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Soybeans Following Corn
<b>Hybrid/Variety</b>	Champion Seed CS288
<b>Application Date/Crop Stage</b>	NA - NA
<b>County</b>	Hamilton

**HERBICIDE TREATMENT**




Yield Data Points

- Extreme
- Untreated
- Non-Trial



ST2008061A

# Trial Grain Yield Map With Soil Survey



ST2008061A

# Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Extreme	Untreated	Extreme	Untreated	
Webster Silty Clay Loam, 0 To 2 Percent Slopes	107	8.0	5.8	67.9	69.0	-1.1
Clarion Loam, 2 To 5 Percent Slopes	138B	9.8	11.3	64.8	66.9	-2.1
Clarion Loam, 2 To 5 Percent Slopes, Moderately Eroded	138B2	4.2	4.3	66.5	67.7	-1.2
Knoke Silty Clay Loam, 0 To 1 Percent Slopes	4	0.5	0.8	68.7	67.1	1.6
Canisteo Silty Clay Loam, 0 To 2 Percent Slopes	507	16.8	19.5	68.9	69.0	-0.1
Nicollet Loam, 1 To 3 Percent Slopes	55	8.1	8.1	68.2	69.4	-1.2
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	0.1	0.1	63.3	64.0	-0.7
Harps Clay Loam, 0 To 2 Percent Slopes	95	1.4	1.2	68.6	69.8	-1.2
<b>YIELD AVERAGE FOR TRIAL</b>				<b>67.6</b>	<b>68.5</b>	<b>-0.9</b>

