



ST2008061B

2008 Aerial Imagery With Soil Survey



ST2008061B

Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics

Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	Champion Seed CS288
Application Date/Crop Stage	8/6/2008 - R3 Beg. Pod
County	Hamilton

FUNGICIDE TREATMENT




Yield Data Points

- Headline
- Untreated
- Non-Trial



ST2008061B

Trial Grain Yield Map With Soil Survey



ST2008061B

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Silty Clay Loam, 0 To 2 Percent Slopes	107	4.9	10.8	69.1	68.0	1.1
Clarion Loam, 2 To 5 Percent Slopes	138B	11.2	12.3	66.5	65.2	1.3
Clarion Loam, 2 To 5 Percent Slopes, Moderately Eroded	138B2	3.6	3.7	69.0	65.2	3.8
Canisteo Silty Clay Loam, 0 To 2 Percent Slopes	507	19.4	15.3	70.3	67.6	2.7
Nicollet Loam, 1 To 3 Percent Slopes	55	7.9	8.8	70.0	67.8	2.2
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	0.0	0.2	0.0	63.5	0.0
Harps Clay Loam, 0 To 2 Percent Slopes	95	1.7	0.2	70.9	65.4	5.5
YIELD AVERAGE FOR TRIAL				69.2	66.9	2.3

