



ST2008375A

# 2008 Aerial Imagery With Soil Survey



ST2008375A

# Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics




<b>Trial Type</b>	Stratego Fungicide
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Soybeans Following Corn
<b>Hybrid/Variety</b>	Liberty Link Foundation Seed
<b>Application Date/Crop Stage</b>	7/15/2008 - R2 Full Bloom
<b>County</b>	Hancock

<b>FUNGICIDE TREATMENT</b>	
Yield Data Points	
●	Stratego
●	Untreated
●	Non-Trial



ST2008375A

# Trial Grain Yield Map With Soil Survey



ST2008375A

# Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Stratego	Untreated	Stratego	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	4.1	5.7	51.7	46.6	5.1
Clarion Loam, 9 To 14 Percent Slopes, Moderately Eroded	138D2	0.0	1.0	0.0	47.4	0.0
Palms Muck, 0 To 1 Percent Slopes	221	8.5	12.3	50.1	48.7	1.4
Spillville Loam, 2 To 5 Percent Slopes	485B	1.0	0.0	54.4	0.0	54.4
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	2.5	3.7	46.0	33.6	12.4
Nicollet Loam, 1 To 3 Percent Slopes	55	8.9	5.4	51.9	43.8	8.1
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	0.0	0.7	0.0	42.1	0.0
Clarion-Sunburg Complex, 5 To 9 Percent Slopes, Moderately Eroded	641C2	0.8	2.0	41.3	49.4	-8.1
Clarion-Sunburg Complex, 9 To 14 Percent Slopes, Moderately Eroded	641D2	3.4	5.7	48.2	39.8	8.4
Okoboji Mucky Silt Loam, 0 To 1 Percent Slopes	90	4.8	6.2	47.8	45.1	2.7
Harps Loam, 0 To 2 Percent Slopes	95	8.6	14.7	50.9	42.9	8.0
<b>YIELD AVERAGE FOR TRIAL</b>				<b>50.1</b>	<b>44.2</b>	<b>5.9</b>

