


ST2008062A

# 2008 Aerial Imagery With Soil Survey



ST2008062A

# Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics

<b>Trial Type</b>	Extreme
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Soybeans Following Corn
<b>Hybrid/Variety</b>	Champion Seed CS288
<b>Application Date/Crop Stage</b>	NA - NA
<b>County</b>	Hardin

**HERBICIDE TREATMENT**



Yield Data Points

- Extreme
- Untreated
- Non-Trial



ST2008062A

# Trial Grain Yield Map With Soil Survey



ST2008062A

## Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Extreme	Untreated	Extreme	Untreated	
Clarion Loam, 2 To 5 Percent Slopes	138B	15.5	19.1	65.1	66.8	-1.7
Webster-Nicollet Complex, 1 To 3 Percent Slopes	329	8.6	12.6	64.9	67.1	-2.2
Canisteo Silty Clay Loam, 0 To 2 Percent Slopes	507	14.0	14.5	58.8	61.1	-2.3
Nicollet Loam, 1 To 3 Percent Slopes	55	5.9	7.5	64.4	66.0	-1.6
Harps Loam, 1 To 3 Percent Slopes	95	0.9	1.3	52.6	58.6	-6.0
<b>YIELD AVERAGE FOR TRIAL</b>				<b>62.8</b>	<b>65.0</b>	<b>-2.2</b>

