




ST2008071A

2008 Aerial Imagery With Soil Survey



ST2008071A


Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics



Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Corn Following Soybeans
Hybrid/Variety	Multiple CSX1078 west 2 reps, CSX1088 east 3 reps
Application Date/Crop Stage	8/4/2008 - R1 Silking
County	Hardin

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008071A

Trial Grain Yield Map With Soil Survey



ST2008071A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Clarion Loam, 2 To 5 Percent Slopes	138B	13.9	14.6	180.8	181.0	-0.2
Clarion Loam, 5 To 9 Percent Slopes, Moderately Eroded	138C2	8.1	6.3	185.9	179.6	6.3
Webster-Nicollet Complex, 1 To 3 Percent Slopes	329	9.7	11.7	153.9	150.0	3.9
Canisteo Silty Clay Loam, 0 To 2 Percent Slopes	507	8.0	8.1	145.9	144.5	1.4
Nicollet Loam, 1 To 3 Percent Slopes	55	3.9	3.8	186.6	165.3	21.3
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	2.2	2.2	134.9	116.6	18.3
Harps Loam, 1 To 3 Percent Slopes	95	3.6	4.0	156.1	140.7	15.4
YIELD AVERAGE FOR TRIAL				167.3	160.7	6.6

