




ST2008071B

# 2008 Aerial Imagery With Soil Survey



ST2008071B

# Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics

<b>Trial Type</b>	Stratego Fungicide
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Corn Following Soybeans
<b>Hybrid/Variety</b>	Multiple CSX1078 west 2 reps, CSX1088 east 3 reps
<b>Application Date/Crop Stage</b>	8/4/2008 - R1 Silking
<b>County</b>	Hardin

**FUNGICIDE TREATMENT**



Yield Data Points

- Stratego
- Untreated
- Non-Trial



ST2008071B

# Trial Grain Yield Map With Soil Survey



ST2008071B

## Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Stratego	Untreated	Stratego	Untreated	
Clarion Loam, 2 To 5 Percent Slopes	138B	13.9	14.2	182.5	181.6	0.9
Clarion Loam, 5 To 9 Percent Slopes, Moderately Eroded	138C2	7.6	6.7	181.1	179.2	1.9
Webster-Nicollet Complex, 1 To 3 Percent Slopes	329	9.8	10.1	143.0	149.0	-6.0
Canisteo Silty Clay Loam, 0 To 2 Percent Slopes	507	8.3	6.7	153.2	144.5	8.7
Nicollet Loam, 1 To 3 Percent Slopes	55	4.1	4.2	172.8	170.6	2.2
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	2.5	3.0	125.9	142.3	-16.4
Harps Loam, 1 To 3 Percent Slopes	95	3.1	5.7	144.5	152.6	-8.1
<b>YIELD AVERAGE FOR TRIAL</b>				<b>163.4</b>	<b>163.4</b>	<b>0.0</b>

