



ST2008076A

2008 Aerial Imagery With Soil Survey



ST2008076A

Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics

Trial Type	Ultimate
Fertilizer Form	NH3
Fertilizer Timing	Sidedress
Treatments	120 lbs N vs 145 lbs N
Previous Crop	Corn
County	Hardin

NITROGEN TREATMENT




Yield Data Points

- 145 lbs N
- 120 lbs N
- Non-Trial



ST2008076A

Trial Grain Yield Map With Soil Survey



ST2008076A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		145 lbs N	120 lbs N	145 lbs N	120 lbs N	
Webster Silty Clay Loam, 0 To 2 Percent Slopes	107	10.2	10.6	160.4	162.5	-2.1
Clarion Loam, 2 To 5 Percent Slopes	138B	5.0	3.9	191.7	171.1	20.6
Nicollet Loam, 1 To 3 Percent Slopes	55	21.7	18.6	176.3	176.9	-0.6
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	2.9	3.6	150.8	121.1	29.7
Harps Loam, 1 To 3 Percent Slopes	95	11.1	12.3	174.5	171.0	3.5
YIELD AVERAGE FOR TRIAL				172.8	167.7	5.1

