



ST2008269B

2008 Aerial Imagery With Soil Survey



ST2008269B

Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics

Trial Type	Stratego & Leverage
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	Croplan Genetics RC2177
Application Date/Crop Stage	8/6/2008 - R3 Beg. Pod
County	Hardin



FUNGICIDE AND INSECTICIDE TREATMENT

- Yield Data Points
- Stratego & Leverage
 - Untreated
 - Non-Trial



ST2008269B

Trial Grain Yield Map With Soil Survey



ST2008269B

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Stratego & Leverage	Untreated	Stratego & Leverage	Untreated	
Clarion Loam, 2 To 5 Percent Slopes	138B	21.2	28.6	69.1	56.7	12.4
Webster-Nicollet Complex, 1 To 3 Percent Slopes	329	9.9	11.0	69.7	56.2	13.5
Canisteo Silty Clay Loam, 0 To 2 Percent Slopes	507	9.6	11.0	58.0	49.5	8.5
Nicollet Loam, 1 To 3 Percent Slopes	55	1.2	2.8	72.1	58.8	13.3
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	0.8	0.9	74.9	55.7	19.2
Harps Loam, 1 To 3 Percent Slopes	95	1.0	1.8	61.3	53.5	7.8
YIELD AVERAGE FOR TRIAL				66.8	55.2	11.6

