



ST2008191A

2008 Aerial Imagery With Soil Survey



ST2008191A

Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics

Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	Latham L-2085
Application Date/Crop Stage	7/28/2008 - R2 Blister
County	Humboldt

FUNGICIDE TREATMENT




Yield Data Points

- Headline
- Untreated
- Non-Trial



ST2008191A

Trial Grain Yield Map With Soil Survey



ST2008191A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Silty Clay Loam, 0 To 2 Percent Slopes	107	40.3	33.6	59.5	58.3	1.2
Clarion Loam, 2 To 5 Percent Slopes	138B	1.1	1.5	59.2	59.0	0.2
Nicollet Loam, 1 To 3 Percent Slopes	55	10.5	9.0	58.0	56.7	1.3
Okoboji Silty Clay Loam, Depressional, 0 To 1 Percent Slopes	6	1.8	2.1	58.4	57.8	0.6
YIELD AVERAGE FOR TRIAL				59.2	58.0	1.2

