



ST2008329A

2008 Aerial Imagery With Soil Survey



ST2008329A


Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics

Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	Latham L-2038
Application Date/Crop Stage	7/25/2008 - R3 Beg. Pod
County	Humboldt

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008329A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Silty Clay Loam, 0 To 2 Percent Slopes	107	6.4	5.4	54.9	53.1	1.8
Clarion Loam, 2 To 5 Percent Slopes	138B	8.7	10.0	51.9	50.2	1.7
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	18.1	17.2	56.3	54.9	1.4
Nicollet Loam, 1 To 3 Percent Slopes	55	16.6	15.9	52.9	51.8	1.1
Depressional, 0 To 1 Percent Slopes	6	0.7	1.0	55.4	53.3	2.1
YIELD AVERAGE FOR TRIAL				54.3	52.7	1.6

