



ST2008339A

# 2008 Aerial Imagery With Soil Survey



ST2008339A


# Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics




<b>Trial Type</b>	Headline Fungicide
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Corn Following Corn (2 Years)
<b>Hybrid/Variety</b>	Agrigold A6325
<b>Application Date/Crop Stage</b>	8/4/2008 - R1 Silking
<b>County</b>	Humboldt

<b>FUNGICIDE TREATMENT</b>	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008339A

# Trial Grain Yield Map With Soil Survey



ST2008339A

# Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Silty Clay Loam, 0 To 2 Percent Slopes	107	22.2	11.8	190.8	189.1	1.7
Clarion Loam, 2 To 5 Percent Slopes	138B	21.4	7.7	202.4	197.9	4.5
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	16.8	9.1	188.4	184.5	3.9
Nicollet Loam, 1 To 3 Percent Slopes	55	6.5	2.3	208.3	194.7	13.6
Okoboji Silty Clay Loam, Depressional, 0 To 1 Percent Slopes	6	1.6	0.5	180.2	177.2	3.0
<b>YIELD AVERAGE FOR TRIAL</b>				<b>195.3</b>	<b>190.2</b>	<b>5.1</b>

