



ST2008188A

2008 Aerial Imagery With Soil Survey



ST2008188A


Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics



Trial Type	Headline Fungicide
Application Method	Aerial Applied
Crop Rotation	Corn Following Soybeans
Hybrid/Variety	Pioneer 35K33
Application Date/Crop Stage	7/30/2008 - R1 Silking
County	Pocahontas

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008188A

Trial Grain Yield Map With Soil Survey



ST2008188A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	7.0	5.5	215.1	202.8	12.3
Clarion Loam, 2 To 5 Percent Slopes	138B	8.0	6.8	213.8	200.0	13.8
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	30.2	25.8	194.9	184.7	10.2
Nicollet Clay Loam, 1 To 3 Percent Slopes	55	4.3	2.7	212.9	202.5	10.4
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	2.6	3.9	224.3	216.7	7.6
Okoboji Mucky Silty Clay Loam, 0 To 1 Percent Slopes	90	1.5	1.7	214.4	209.2	5.2
YIELD AVERAGE FOR TRIAL				203.7	193.7	10.0

