



ST2008188B

2008 Aerial Imagery With Soil Survey



ST2008188B


Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics


Trial Type	Headline Fungicide
Application Method	Aerial Applied
Crop Rotation	Corn Following Soybeans
Hybrid/Variety	Pioneer 36y86
Application Date/Crop Stage	7/30/2008 - R1 Silking
County	Pocahontas

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008188B

Trial Grain Yield Map With Soil Survey



GRAIN YIELD (bu/a)

- 156.2 - 202.6
- 202.6 - 212.9
- 212.9 - 219.9
- 219.9 - 226.0
- 226.0 - 233.9
- 233.9 - 262.6

□ Soil Survey



ST2008188B

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	8.2	12.6	218.5	213.4	5.1
Clarion Loam, 2 To 5 Percent Slopes	138B	5.2	8.9	221.7	221.6	0.1
Clarion Loam, 5 To 9 Percent Slopes, Moderately Eroded	138C2	2.0	4.9	204.0	214.1	-10.1
Knoke Silty Clay Loam, 0 To 1 Percent Slopes	4	2.8	4.3	211.9	203.8	8.1
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	6.2	12.0	220.1	213.3	6.8
Nicollet Clay Loam, 1 To 3 Percent Slopes	55	5.9	7.7	226.0	217.0	9.0
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	0.4	1.2	228.6	217.7	10.9
Okoboji Mucky Silty Clay Loam, 0 To 1 Percent Slopes	90	6.6	11.1	231.7	225.6	6.1
YIELD AVERAGE FOR TRIAL				221.6	216.6	5.0

