



ST2008229B

2008 Aerial Imagery With Soil Survey



ST2008229B

Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics

Trial Type	Stratego Fungicide
Application Method	Aerial Applied
Crop Rotation	Corn Following Soybeans
Hybrid/Variety	Dekalb DKC50-48
Application Date/Crop Stage	7/30/2008 - R1 Silking
County	Pocahontas

FUNGICIDE TREATMENT



Yield Data Points

- Stratego
- Untreated
- Non-Trial



ST2008229B

Trial Grain Yield Map With Soil Survey



ST2008229B

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Stratego	Untreated	Stratego	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	24.2	26.0	208.2	208.3	-0.1
Clarion Loam, 2 To 5 Percent Slopes	138B	3.2	3.6	210.8	209.1	1.7
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	5.6	4.1	211.5	209.7	1.8
Nicollet Clay Loam, 1 To 3 Percent Slopes	55	15.8	15.5	207.8	207.4	0.4
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	1.1	1.0	209.6	205.5	4.1
YIELD AVERAGE FOR TRIAL				208.6	208.1	0.5

