




ST2008185A

# 2008 Aerial Imagery With Soil Survey



ST2008185A

# Treatment Type and Location of Trial Within Field




<b>Trial Type</b>	Headline Fungicide
<b>Application Method</b>	Ground Applied
<b>Crop Rotation</b>	Soybeans Following Corn
<b>Hybrid/Variety</b>	Pioneer 92M54
<b>Application Date/Crop Stage</b>	8/7/2008 - R3 Beg. Pod
<b>County</b>	Polk

**FUNGICIDE TREATMENT**




Yield Data Points

- Headline
- Untreated
- Non-Trial



ST2008185A

# Trial Grain Yield Map With Soil Survey



ST2008185A

## Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Clarion Loam, Moderately Coarse Substratum, 2 To 5 Percent Slopes	138B	7.3	10.6	55.0	53.0	2.0
Canisteo Clay Loam, Moderately Coarse Substratum, 0 To 2 Percent Slopes	507	22.9	32.2	52.1	49.2	2.9
Nicollet Loam, 1 To 3 Percent Slopes	55	11.2	15.8	54.0	50.4	3.6
<b>YIELD AVERAGE FOR TRIAL</b>				<b>53.1</b>	<b>50.2</b>	<b>2.9</b>

