




ST2008169A

2008 Aerial Imagery With Soil Survey



ST2008169A


Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics




Trial Type	Headline Fungicide
Application Method	Aerial Applied
Crop Rotation	Corn Following Soybeans
Hybrid/Variety	Pioneer 33M16
Application Date/Crop Stage	8/19/2008 - R2 Blister
County	Story

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008169A

Trial Grain Yield Map With Soil Survey



ST2008169A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	2.7	4.3	230.5	224.9	5.6
Clarion Loam, 2 To 5 Percent Slopes	138B	22.6	23.7	240.0	230.7	9.3
Clarion Loam, 5 To 9 Percent Slopes, Moderately Eroded	138C2	1.0	2.4	250.4	250.6	-0.2
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	7.4	8.2	237.3	240.1	-2.8
Nicollet Loam, 1 To 3 Percent Slopes	55	11.3	14.6	241.9	233.7	8.2
Harps Loam, 1 To 3 Percent Slopes	95	0.7	1.2	253.0	264.4	-11.4
YIELD AVERAGE FOR TRIAL				239.9	234.1	5.8

