




ST2008170A

# 2008 Aerial Imagery With Soil Survey



ST2008170A

# Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics






<b>Trial Type</b>	Headline Fungicide
<b>Application Method</b>	Aerial Applied
<b>Crop Rotation</b>	Corn Following Corn (2 Years)
<b>Hybrid/Variety</b>	Pioneer 33D14
<b>Application Date/Crop Stage</b>	8/19/2008 - R2 Blister
<b>County</b>	Story

<b>FUNGICIDE TREATMENT</b>	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008170A

# Trial Grain Yield Map With Soil Survey



ST2008170A

## Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	5.7	14.2	175.1	174.8	0.3
Coland Clay Loam, 0 To 2 Percent Slopes	135	0.0	0.2	0.0	182.8	0.0
Clarion Loam, 2 To 5 Percent Slopes	138B	14.6	28.0	171.6	172.8	-1.2
Clarion Loam, 5 To 9 Percent Slopes, Moderately Eroded	138C2	5.7	10.2	181.9	186.9	-5.0
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	1.1	3.1	184.4	175.9	8.5
Nicollet Loam, 1 To 3 Percent Slopes	55	3.9	8.8	171.8	169.5	2.3
Zenor Sandy Loam, 5 To 9 Percent Slopes, Moderately Eroded	828C2	1.1	3.4	154.6	170.8	-16.2
<b>YIELD AVERAGE FOR TRIAL</b>				<b>173.9</b>	<b>175.0</b>	<b>-1.1</b>

