




ST2008177A

# 2008 Aerial Imagery With Soil Survey



ST2008177A


# Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)  
Are Not Used In Summary Statistics




<b>Trial Type</b>	Headline Fungicide
<b>Application Method</b>	Aerial Applied
<b>Crop Rotation</b>	Corn Following Corn (2 Years)
<b>Hybrid/Variety</b>	Dekalb DKC62-29
<b>Application Date/Crop Stage</b>	8/19/2008 - R2 Blister
<b>County</b>	Story

<b>FUNGICIDE TREATMENT</b>	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008177A

# Trial Grain Yield Map With Soil Survey



ST2008177A

## Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	12.5	10.8	197.5	180.2	17.3
Clarion Loam, 2 To 5 Percent Slopes	138B	12.3	14.6	203.9	194.6	9.3
Canisteo Clay Loam, 0 To 2 Percent Slopes	507	5.7	4.2	199.7	190.3	9.4
Nicollet Loam, 1 To 3 Percent Slopes	55	18.9	17.5	201.5	186.3	15.2
Harps Loam, 1 To 3 Percent Slopes	95	1.8	1.6	200.3	191.2	9.1
<b>YIELD AVERAGE FOR TRIAL</b>				<b>200.9</b>	<b>188.0</b>	<b>12.9</b>

