




ST2008178A

2008 Aerial Imagery With Soil Survey



ST2008178A


Treatment Type and Location of Trial Within Field



Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics




Trial Type	Headline Fungicide
Application Method	Ground Applied
Crop Rotation	Soybeans Following Corn
Hybrid/Variety	Prairie Brand PB2557
Application Date/Crop Stage	8/6/2008 - R3 Beg. Pod
County	Story

FUNGICIDE TREATMENT	
Yield Data Points	
●	Headline
●	Untreated
●	Non-Trial



ST2008178A

Trial Grain Yield Map With Soil Survey



ST2008178A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Webster Clay Loam, 0 To 2 Percent Slopes	107	5.5	6.1	54.8	50.9	3.9
Clarion Loam, 2 To 5 Percent Slopes	138B	30.0	28.7	57.6	54.3	3.3
Clarion Loam, 5 To 9 Percent Slopes, Moderately Eroded	138C2	4.1	3.8	57.0	51.2	5.8
Dickinson Fine Sandy Loam, 2 To 5 Percent Slopes	175B	0.0	1.2	0.0	62.1	0.0
Farrar Fine Sandy Loam, 2 To 5 Percent Slopes	253B	1.6	2.1	50.5	51.0	-0.5
Nicollet Loam, 1 To 3 Percent Slopes	55	7.4	7.3	52.8	53.9	-1.1
Harps Loam, 1 To 3 Percent Slopes	95	1.6	0.4	53.4	53.8	-0.4
YIELD AVERAGE FOR TRIAL				56.2	53.6	2.6

