



ST2008187A

2008 Aerial Imagery With Soil Survey



ST2008187A

Treatment Type and Location of Trial Within Field




Trial Type	Headline Fungicide
Application Method	Aerial Applied
Crop Rotation	Corn Following Soybeans
Hybrid/Variety	Pioneer 33M16
Application Date/Crop Stage	8/19/2008 - R2 Blister
County	Story

FUNGICIDE TREATMENT




Yield Data Points

- Headline
- Untreated
- Non-Trial



ST2008187A

Trial Grain Yield Map With Soil Survey



ST2008187A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		Headline	Untreated	Headline	Untreated	
Clarion Loam, 2 To 5 Percent Slopes	138B	18.7	39.6	242.6	234.5	8.1
Clarion Loam, 5 To 9 Percent Slopes, Moderately Eroded	138C2	1.4	4.8	218.9	211.7	7.2
Clarion Loam, 9 To 14 Percent Slopes, Moderately Eroded	138D2	2.3	9.0	213.2	196.0	17.2
Nicollet Loam, 1 To 3 Percent Slopes	55	3.7	6.5	251.6	241.9	9.7
Storden Loam, 14 To 18 Percent Slopes, Severely Eroded	62E3	0.4	0.9	236.4	249.6	-13.2
Clarion-Storden Loams, 9 To 14 Percent Slopes, Moderately Erode	638D2	4.9	7.9	204.0	216.5	-12.5
YIELD AVERAGE FOR TRIAL				234.4	226.7	7.7

