



ST2008041A

2008 Aerial Imagery With Soil Survey



ST2008041A

Treatment Type and Location of Trial Within Field




Non-Trial Yield Points (in grey)
Are Not Used In Summary Statistics






Trial Type	75/125
Fertilizer Form	NH3
Fertilizer Timing	Spring
Treatments	75 lbs N vs 125 lbs N
Previous Crop	Soybeans
County	Wright

NITROGEN TREATMENT	
Yield Data Points	
●	125 lbs N
●	75 lbs N
●	Non-Trial



ST2008041A

Trial Grain Yield Map With Soil Survey



ST2008041A

Summary Statistics By Soil Type and By Individual Treatment Strip

Soil Name	Label	Percent of Trial		Yield (bu/a)		Yield Difference
		125 lbs N	75 lbs N	125 lbs N	75 lbs N	
Brownton Silty Clay Loam, 0 To 2 Percent Slopes	1507	32.0	27.1	194.7	177.0	17.7
Ottosen Clay Loam, 1 To 3 Percent Slopes	288	7.5	6.8	195.8	179.4	16.4
Okoboji Silty Clay Loam, 0 To 1 Percent Slopes	6	1.1	0.6	162.2	167.8	-5.6
Okoboji Mucky Silty Clay Loam, 0 To 1 Percent Slopes	90	0.1	0.0	140.9	165.1	-24.2
Harps Clay Loam, 0 To 2 Percent Slopes	95	14.0	10.7	193.4	171.5	21.9
YIELD AVERAGE FOR TRIAL				193.7	175.9	17.8

